Presented at Flavor Physics and CP Violation (FPCP 2012), Hefei, China, May 21-25, 2012

arXiv:1208.4980v1 [hep-ex] 24 Aug 2012

CP Violation in Other B_s^0 Decays

L. Zhang (On behalf of LHCb Collaboration)
Syracuse University, Syracuse, New York, 13244, USA

The recent experimental results of CP violation in B_s^0 decays other than in the $J/\psi\phi$ final state are discussed. Included are the resonant components and ϕ_s determination in $B_s^0 \to J/\psi\pi^+\pi^-$, CP asymmetries in $B_s^0 \to h^+h^{(\prime)-}$ decays, and the B_s^0 effective lifetimes in the CP-even state K^+K^- and the CP-odd state $J/\psi f_0(980)$.

1. Introduction

The time evolution of the B_q^0-$\overline{B}{}_q^0$ system is described by the Schrödinger equation

$$i\frac{\partial}{\partial t}\begin{pmatrix}|B_q^0(t)\rangle\\ |\overline{B}{}_q^0(t)\rangle\end{pmatrix} = \left(\mathbf{M}^q - \frac{i}{2}\mathbf{\Gamma}^q\right)\begin{pmatrix}|B_q^0(t)\rangle\\ |\overline{B}{}_q^0(t)\rangle\end{pmatrix}, \tag{1}$$

where the $\mathbf{M}^q$ and $\mathbf{\Gamma}^q$ matrices are Hermitian, and CPT invariance requires $M_{11}^q = M_{22}^q$ and $\Gamma_{11}^q = \Gamma_{22}^q$. The off-diagonal elements, M_{12}^q and Γ_{12}^q, of these matrices describe the off-shell (dispersive) and on-shell (absorptive) contributions to B_q^0-$\overline{B}{}_q^0$ mixing, respectively.

The mixing can be described by three physical quantities: $|M_{12}^q|$, $|\Gamma_{12}^q|$ and the relative phase $\phi_{12}^q = \arg\left(-\frac{M_{12}^q}{\Gamma_{12}^q}\right)$. They are related to the experimental observables [Lenz 2012] as: $\Delta M_q \simeq 2|M_{12}^q|$, $\Delta\Gamma_q \simeq 2|\Gamma_{12}^q|\cos\phi_{12}^q$, and $a_{\rm sl}^q \simeq \frac{\Delta\Gamma_q}{\Delta m_q}\tan\phi_{12}^q$, where ΔM_q is the mass difference between the heavy and the light mass eigenstates, $-\Delta\Gamma_q$ the width difference, and $a_{\rm sl}^q$ the semileptonic (or flavor specific) CP asymmetry. The phase difference, denoted as ϕ_q, between $\arg(M_{12}^q)$ and the decay phase in $b\to c\bar{c}s$ transitions is also an experimental observable. In the B_s case, $\phi_s = -2\arg[V_{ts}V_{tb}^*/V_{cs}V_{cb}^*]$, is small and accurately predicted as $-0.0363^{+0.0016}_{-0.0015}$ rad [Charles 2011]. New Physics [Bobeth and Haisch 2011] in mixing could add new phases to M_{12}^q or/and Γ_{12}^q and modify ϕ_{12}^q or/and ϕ_q from their Standard Model (SM) predictions.

2. Formalism

The time dependent decay rates for initial B_q^0 or $\overline{B}{}_q^0$ decays to a CP eigenstate ($f_{\rm CP}$) are:

$$\Gamma(B_q^0(t)\to f_{\rm CP}) = \mathcal{N}e^{-\Gamma_q t}\left\{\frac{1+|\lambda|^2}{2}\cosh\frac{\Delta\Gamma_q t}{2} + \frac{1-|\lambda|^2}{2}\cos(\Delta m_q t) - {\rm Re}(\lambda)\sinh\frac{\Delta\Gamma_q t}{2} - {\rm Im}(\lambda)\sin(\Delta m_q t)\right\},$$

$$\Gamma(\overline{B}{}_q^0(t)\to f_{\rm CP}) = \mathcal{N}e^{-\Gamma_q t}\left\{\frac{1+|\lambda|^2}{2}\cosh\frac{\Delta\Gamma_q t}{2} - \frac{1-|\lambda|^2}{2}\cos(\Delta m_q t) - {\rm Re}(\lambda)\sinh\frac{\Delta\Gamma_q t}{2} + {\rm Im}(\lambda)\sin(\Delta m_q t)\right\}, \tag{2}$$

where $\lambda = (q/p)(\bar{A}_f/A_f)$, Γ_q is average width of two mass eigenstates, A_f ($\bar{A}_f$) is the amplitude of B_q^0 ($\overline{B}{}_q^0$) decay, and $|p/q| = 1$ is used assuming no CP violation in the mixing.

The CP asymmetry is

$$A_{f_{CP}}(t) \equiv \frac{\Gamma(\overline{B}{}_q^0(t)\to f_{CP}) - \Gamma(B_q^0(t)\to f_{CP})}{\Gamma(\overline{B}{}_q^0(t)\to f_{CP}) + \Gamma(B_q^0(t)\to f_{CP})} = \frac{A^{\rm dir}\cos(\Delta m_q t) + A^{\rm mix}\sin(\Delta m_q t)}{\cosh\frac{\Delta\Gamma_q t}{2} + A^{\Delta\Gamma}\sinh\frac{\Delta\Gamma_q t}{2}}. \tag{3}$$

There are three CP variables but only two are independent: direct CP asymmetry $A^{\rm dir} = \frac{|\lambda|^2-1}{|\lambda|^2+1}$, mixing induced CP asymmetry $A^{\rm mix} = \frac{2{\rm Im}\lambda}{|\lambda|^2+1}$ and $A^{\Delta\Gamma} = -\frac{2{\rm Re}\lambda}{|\lambda|^2+1}$ satisfying $(A^{\rm dir})^2 + (A^{\rm mix})^2 + (A^{\Delta\Gamma})^2 = 1$.

3. Resonant components and ϕ_s determination in $B_s^0 \to J/\psi\pi^+\pi^-$

Motivated by a predication in Ref. [Stone and Zhang 2009], the LHCb collaboration made the first observation of $B_s^0 \to J/\psi f_0(980), f_0(980) \to \pi^+\pi^-$ [Aaij 2011], which was subsequently confirmed by others [Li 2011, Aaltonen 2011a, Abazov 2012a]. This mode is a CP-odd eigenstate and can be used to determine ϕ_s without the need for an angular analysis, as is required in the $J/\psi\phi$ final state [Aaij 2012a, Abazov 2012b, Aaltonen 2011b]. With 0.4 fb^{-1} data, LHCb used the candidates within ± 90 MeV of $f_0(980)$ mass and measured $\phi_s = -0.44 \pm 0.44 \pm 0.02$ rad [Aaij 2012b]. Whenever two uncertainties are given, the first is statistical and the second systematic. However, the used events are only about half of $J/\psi\pi^+\pi^-$ signal.

To optimize $J/\psi\pi^+\pi^-$ usefulness, we need to understand the CP content of this final state. With 1.0 fb^{-1} data, LHCb [Aaij 2012c] preformed a modified Dalitz-plot analysis which fits to the $m^2(\pi^+\pi^-)$, $m^2(J/\psi\pi^+)$ and $J/\psi \to \mu^+\mu^-$ helicity angle ($\theta_{J/\psi}$) distributions after integrating the angle between J/ψ and $\pi^+\pi^-$ decay planes. Different from the classical "Dalitz-plot" analysis, the vector J/ψ particle has 3 helicity amplitudes that must be considered. The projection of $m^2(\pi^+\pi^-)$ overlayed with the best fit is shown in Fig. 1. The components and fractions of the best fit are given in Table I. The fraction of $J/\psi\rho$ is measured less than 1.6% at 95% confidence level (CL); the $J/\psi\rho$ final state only can be present in higher order processes. The final states listed in Table I are all CP-odd states, expect for $f_2(1270)$ with helicity $|\Lambda| = 1$ which has mixed-CP. Combining with this and ρ, the fraction of CP-even states is less than 2.3% at 95% CL. So the whole mode is dominated by CP-odd state, and can be used for ϕ_s measurement without need of angular analysis. The relative branching ratio between $B_s^0 \to J/\psi\pi^+\pi^-$ and $J/\psi\phi$ is measured as (19.79±0.47±0.52)% [Aaij 2012c].



Figure 1: Dalitz fit project of $m^2(\pi^+\pi^-)$ from the best fit. The points with error bars are data, the signal fit is shown with a (red) dashed line, the background with a (black) dotted line, and the (blue) solid line represents the total.

The $B_s^0 \to J/\psi\pi^+\pi^-$ decay complements $J/\psi\phi$ in determining ϕ_s. The event yield is about 40% of $J/\psi\phi(\phi \to K^+K^-)$. In B_s^0 decay to a CP-odd final state via $b \to c\bar{c}s$ transition, the time dependent decay rates are given in Eq. (2) with $\lambda = -e^{-\phi_s}$. We also need to take into account experimental effects, such as acceptance in decay time, time resolution, and dilution due to wrong flavor tagging. The decay time resolution in LHCb is about 40 fs thanks to the excellent vertex detector and large momentum of B_s^0.

LHCb updated the ϕ_s measurement using an sample of 1 fb^{-1} data [Aaij 2012d]. The $m(J/\psi\pi^+\pi^-)$ and $m(\pi^+\pi^-)$ distributions are shown in Fig. 2 after applying a Boosted Decision Tree selection rather than a cut-based selection used in Fig. 1 . With about 7400 B_s^0 signal candidates, LHCb finds $\phi_s = -0.019^{+0.173+0.004}_{-0.174-0.003}$ rad, consistent with the SM expectation [Charles 2011]. The largest systematic uncertainty arises from allowing direct CP violation.

Table I: Resonance fractions in $B_s^0 \to J/\psi\pi^+\pi^-$ over the full mass range [Aaij 2012c]. The final-state helicity of the D-wave is denoted by Λ. Only statistical uncertainties are quoted.

Resonance	Normalized fraction (%)
$f_0(980)$	69.7 ± 2.3
$f_0(1370)$	21.2 ± 2.7
non-resonant $\pi^+\pi^-$	8.4 ± 1.5
$f_2(1270)$, $\Lambda = 0$	0.49 ± 0.16
$f_2(1270)$, $\lvert\Lambda\rvert = 1$	0.21 ± 0.65

Here ϕ_s changes by -0.0020 from the default value and $|\lambda| = 0.89 \pm 0.13$ consistent with no direct CP violation. The systematic uncertainty due to a possible CP-even component is -0.0008. The measured ϕ_s is consistent with the LHCb preliminary result $-0.001 \pm 0.101 \pm 0.027$ rad using $B_s^0 \to J/\psi\phi$ [Aaij 2012e]. Combining the two LHCb measurements gives $\phi_s = -0.002 \pm 0.083 \pm 0.027$ rad [Aaij 2012e].



Figure 2: Left: The $m(J/\psi\pi^+\pi^-)$ distribution from the candidates with $m(\pi^+\pi^-) \in [775, 1550]$ MeV, shown also the B_s^0 signal as the red solid line and other various background components. Right: The $m(\pi^+\pi^-)$ from the candidates with ±20 MeV around B_s^0 mass peak. The arrows indicate the region used for ϕ_s measurement and the red dash line shows the background.

Other possible modes for measuring ϕ_s are

- $B_s^0 \to J/\psi f_2'(1525)$ which was first observed by LHCb [Aaij 2012f]. This mode could be useful but it needs to include additional D-wave in transversity amplitudes. Sizeable S-wave over the entire $m(K^+K^-)$ in $J/\psi K^+K^-$ is also seen [Aaij 2012f].
- $B_s^0 \to J/\psi\eta^{(\prime)}$ which were first observed by Belle [Li 2012]. They are CP-even states with large branching fraction, but neutral detection is difficult for experiments in a hadron collider.
- $B_s^0 \to \psi(2S)\phi, \psi(2S) \to \mu^+\mu^-$ and $B_s^0 \to D_s^+D_s^-$. The event yields are of order of 5-10% of $(J/\psi \to \mu^+\mu^-)(\phi \to K^+K^-)$.

4. CP asymmetries in $B_s^0 \to h^+h^{(\prime)-}$ decays

Two-body charmless B_s^0 decays have significant contribution of penguin diagrams, providing an entry point for New Physics. Under U-spin symmetry, where s quarks are changed to d quarks, $B_s^0 \to K^+K^-$ is analogous to $B^0 \to K^+\pi^-$ and $B_s^0 \to K^-\pi^+$ to $B^0 \to \pi^+\pi^-$. Specifically it turns out that $A_{KK}^{\rm dir}$ in B_s^0 should be equal to the



Figure 3: Invariant $K\pi$ mass spectra for (a) $K^+\pi^-$ and (b) $K^-\pi^+$ combinations.

direct CP asymmetry in $B^0 \to K^+\pi^-$, in the limit of exact U-spin symmetry and neglecting certain diagrams with annihilation topologies contributing to the decay amplitude [Fleischer 1999].

The direct CP asymmetry in the flavor-specific decay $B_s^0 \to (f = K^-\pi^+)$,

$$A_{\mathrm{CP}}(B_s^0 \to K\pi) = \frac{|\overline{A}_{\bar{f}}|^2 - |A_f|^2}{|\overline{A}_{\bar{f}}|^2 + |A_f|^2}, \tag{4}$$

is measured by LHCb using 0.35 fb^{-1} data, where an untagged time-integrated method is used. The measured charge asymmetry after correcting detection and production biases is equal to $A_{\mathrm{CP}} + \mathcal{O}(a_{\mathrm{sl}}^s)$. The semileptonic asymmetry is measured as $a_{\mathrm{sl}}^s = (-1.81 \pm 1.06)\%$ by the D0 Collaboration [Abazov 2011], where the uncertainty is sum of statistical and systematic. The LHCb Collaboration recently measured $a_{\mathrm{sl}}^s = (-0.24 \pm 0.54 \pm 0.33)\%$ [Aaij 2012g]. Thus $\mathcal{O}(a_{\mathrm{sl}}^s)$ is negligible compared to the measured CP asymmetry shown later. The $K^+\pi^-$ and $K^-\pi^+$ mass spectra for the candidates are shown in Fig. 3. A Clear difference in the yields of two peaks at the B_s^0 mass is seen. LHCb finds $A_{\mathrm{CP}}(B_s^0 \to K\pi) = 0.27 \pm 0.08 \pm 0.02$ [Aaij 2012h]. It is 3.3σ from zero, providing the first evidence for CP violation in the decays of B_s^0 mesons. The result for $A_{\mathrm{CP}}(B_s^0 \to K\pi)$ is consistent with the only previous measurement from CDF of $0.39 \pm 0.15 \pm 0.08$ [Aaltonen 2011c].

LHCb [Aaij 2012i] made the first measurement of CP asymmetry in $B_s^0 \to K^+K^-$. It's a time-dependent flavor-tagged analysis. The large statistical sample $B^0 \to K^-\pi^+$ is used to measure flavor tagging efficiency and mistag probability that are then constrained in the fit to $B_s^0 \to K^+K^-$. The mass distribution of selected candidates is shown in Fig. 4 (left); fit obtains 7155 ± 97 signal candidates. The measured CP asymmetries (preliminary) are

$$\begin{aligned} A_{KK}^{\mathrm{dir}} &= 0.02 \pm 0.18 \pm 0.04, \\ A_{KK}^{\mathrm{mix}} &= 0.17 \pm 0.18 \pm 0.05. \end{aligned} \tag{5}$$

The raw asymmetry is shown in Fig. 4 (right).

5. Effective lifetimes of B_s^0 from CP-eigenstate final states

The untagged decay time distribution summing B_s^0 and $\overline{B}_s^0$ is

$$\Gamma(t) \propto (1 - A^{\Delta\Gamma_s})e^{-\Gamma_L t} + (1 + A^{\Delta\Gamma_s})e^{-\Gamma_H t}, \tag{6}$$

where $A^{\Delta\Gamma_s} = -\dfrac{2\mathrm{Re}\lambda}{|\lambda|^2 + 1}$, as in Eq. (2), and τ_H and τ_L are the lifetimes of heavy and light mass eigenstates. The CP-odd state $J/\psi f_0(980)$ has $A^{\Delta\Gamma_s} = \cos\phi_s = 1$ and CP-even state K^+K^- has $A^{\Delta\Gamma_s} \approx -1$ (the small difference from unity is discussed in Ref. [Fleischer and Knegjens 2011]). Therefore, their effective lifetimes are approximately equal to the lifetimes of heavy and light mass eigenstates, respectively.



Figure 4: Left: Invariant mass projection for the $B_s^0 \to K^+K^-$. The data are described by the overall fit result (solid blue line) which is the sum of: $B_s^0 \to K^+K^-$ signal (dotted dark yellow line); $B^0 \to K\pi$ cross-feed background (dashed double-dotted red line); partially reconstructed three-body (dashed orange line) and combinatorial (dashed dotted grey line) backgrounds. Right: Time-dependent raw asymmetry in the $B_s^0 \to K^+K^-$ signal mass region with the result of the fit superimposed. The offset $t_0 = 0.95$ ps corresponds to the selection requirement where the decay time acceptance starts.

The effective lifetime in $J/\psi f_0(980)$ final state is measured to be $1.700 \pm 0.040 \pm 0.026$ ps by LHCb [Aaij 2012j] and $1.70^{+0.12}_{-0.11} \pm 0.03$ ps by CDF [Aaltonen 2011a]. The most precise measurement on the effective lifetime in K^+K^- final state comes from LHCb. The measured value is $1.455 \pm 0.046 \pm 0.006$ ps [Aaij 2012k]. Those measurements can be used to improve the determination of Γ_s and $\Delta\Gamma_s$ using the method carried out by HFAG [Amhis 2012].

6. Summary

In summary, the latest world best measurements on CP violation in other B_s^0 decays are discussed. They are mostly provided by LHCb collaboration thanks to the larger statics and better detector performance.

Acknowledgments

Work supported by U.S. National Science Foundation.

References

A. Lenz (2012), arXiv:1205.1444.

J. Charles, O. Deschamps, S. Descotes-Genon, R. Itoh, H. Lacker, et al., Phys. Rev. **D84**, 033005 (2011), arXiv:1106.4041.

C. Bobeth and U. Haisch (2011), arXiv:1109.1826.

S. Stone and L. Zhang, Phys. Rev. **D79**, 074024 (2009), arXiv:0812.2832.

R. Aaij et al. (LHCb collaboration), Phys. Lett. **B698**, 115 (2011), arXiv:1102.0206.

J. Li et al. (Belle collaboration), Phys. Rev. Lett. **106**, 121802 (2011), arXiv:1102.2759.

T. Aaltonen et al. (CDF collaboration), Phys. Rev. **D84**, 052012 (2011a), arXiv:1106.3682.

V. M. Abazov et al. (D0 collaboration), Phys.Rev. **D85**, 011103 (2012a), arXiv:1110.4272.

R. Aaij et al. (LHCb collaboration), Phys. Rev. Lett. **108**, 101803 (2012a), arXiv:1112.3183.

V. M. Abazov et al. (D0 collaboration), Phys.Rev. **D85**, 032006 (2012b), arXiv:1109.3166.

T. Aaltonen et al. (CDF collaboration) (2011b), arXiv:1112.1726.

R. Aaij et al. (LHCb collaboration), Phys. Lett. **B707**, 497 (2012b), arXiv:1112.3056.

R. Aaij et al. (LHCb collaboration) (2012c), arXiv:1204.5643.

R. Aaij et al. (LHCb collaboration), Phys.Lett. **B713**, 378 (2012d), arXiv:1204.5675.

R. Aaij et al. (LHCb collaboration) (2012e), LHCb-CONF-2012-002.

R. Aaij et al. (LHCb Collaboration), Phys.Rev.Lett. **108**, 151801 (2012f), arXiv:1112.4695.

J. Li et al. (Belle Collaboration), Phys.Rev.Lett. **108**, 181808 (2012), arXiv:1202.0103.

R. Fleischer, Phys.Lett. **B459**, 306 (1999), arXiv:hep-ph/9903456.

V. M. Abazov et al. (D0 Collaboration), Phys.Rev. **D84**, 052007 (2011), 1106.6308.

R. Aaij et al. (LHCb collaboration) (2012g), LHCb-CONF-2012-022.

R. Aaij et al. (LHCb Collaboration), Phys.Rev.Lett. **108**, 201601 (2012h), arXiv:1202.6251.

T. Aaltonen et al. (CDF Collaboration), Phys.Rev.Lett. **106**, 181802 (2011c), arXiv:1103.5762.

R. Aaij et al. (LHCb collaboration) (2012i), LHCb-CONF-2012-007.

R. Fleischer and R. Knegjens, Eur.Phys.J. **C71**, 1789 (2011), arXiv:1109.5115.

R. Aaij et al. (LHCb collaboration) (2012j), arXiv:1207.0878.

R. Aaij et al. (LHCb collaboration) (2012k), arXiv:1207.5993.

Y. Amhis et al. (Heavy Flavor Averaging Group) (2012), arXiv:1207.1158.